



ORIGINAL ARTICLE

Effects of polyphenolic antioxidants on exercise-induced oxidative stress

J.M. Morillas-Ruiz^{a,*}, J.A. Villegas García^b, F.J. López^b,
M.L. Vidal-Guevara^c, P. Zafrilla^a

^a*E.U. Human and Dietetic Nutrition, San Antonio Catholic University, Campus Los Jerónimos, s/n. 30107, Murcia*

^b*U.C. Physiology of the Exercise. San Antonio Catholic University, Murcia*

^c*Research and Development Department, Hero Spain, Murcia*

Received 1 August 2005; accepted 4 November 2005

KEYWORDS

Polyphenols;
Antioxidants;
Oxidative stress;
Exercise;
Lipid and protein
peroxidation

Summary Polyphenols are of increasing interest to consumers and food manufacturers for several reasons. Commonly referred to as antioxidants (they are the most abundant antioxidants in our diets), they may prevent various oxidative stress-related diseases, such as cancer, cardiovascular disease, inflammation and others. Physical activity is known to induce oxidative stress in individuals after intensive exercise. In this study, the effect of the flavonoid contents (which are the most abundant polyphenols) was investigated, as the only antioxidant in a replacement drink designed for sportsmen on various oxidative stress biomarkers after two identical trials of sub-maximal aerobic exercise, in a group of 30 sportsmen. In one of the trials, the cyclists consumed the antioxidant supplement (with 2.3 g polyphenols/trial), and in another they consumed a placebo. Blood samples were collected both at rest and after exercise immediately and 45 minutes (min) later, for measurements of plasmatic indices of oxidative stress: lipid oxidation (TBARS), total antioxidant status (TAS); protein oxidation (carbonyl groups, CO) and the lactate dehydrogenase (LDH) and creatine kinase (CK) enzymes for each trial. All values were adjusted for changes in plasma volume. No changes were detected in plasma TAS and LDH after exercise or after the polyphenolic supplement. CK and TBARS increased after exercise in both tests. However, in response to strenuous exercise, the polyphenol-supplemented test showed a smaller increase in plasma TBARS and CK than the placebo test. CO increased by 12% in response to the placebo test, whereas it decreased by 23% in the polyphenol-supplement test. This may indicate that the antioxidant supplement offered protection against exercise-induced oxidative stress.

© 2005 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

*Corresponding author. Tel.: +34 968 278753; fax: +34 968 278726.
E-mail address: jmmorillas@pdi.ucam.edu (J.M. Morillas-Ruiz).

Introduction

Flavonoids are polyphenolic compounds found in rich abundance in all land plants.¹ Owing to their polyphenolic nature, flavonoids often exhibit strong antioxidant properties, akin to α -tocopherol, which they structurally resemble and can replace in some model systems.^{1,2} Although animals do not directly synthesize flavonoids, their diet contains them in large amounts. One estimate has the average daily consumption of flavonoids by humans as 1 g, an amount far greater than that of other dietary antioxidants such as ascorbate or α -tocopherol.³ Given the prevalence of these substances in our diet, and their demonstrable antioxidant activity, it is only reasonable to suppose that animals have evolved the capacity to take advantage of the beneficial properties of flavonoids. Bioavailability studies have shown that the concentrations of intact flavonoids in human plasma rarely exceed 1 μ M when the quantities of polyphenols ingested do not exceed those commonly ingested with our diets. These maximum concentrations are most often reached 1–2 h after ingestion^{4,5} except for polyphenols, which are absorbed only after partial degradation by the microflora in the colon. For most flavonoids absorbed in the small intestine, the plasmatic concentration decreases rapidly (half-life period of 1–2 h).

During strenuous exercise, there is a dramatic increase in oxygen uptake in various organs, particularly in the skeletal muscle.⁶ Oxygen-centered radicals are produced in intermediate metabolism. The resting body is equipped with both non-enzymatic and enzymatic antioxidant reserves to prevent the potential harmful effects of reactive oxygen species (ROS).⁷ The fine physiological balance between oxidative reactions and antioxidant capacity may be perturbed by intense physical activity. ROS released causes lipid peroxidation of polyunsaturated fatty acids in biological membranes and blood, disturbing cell functions.⁸ Malondialdehyde (MDA), a by-product of lipid peroxide, is the most frequently studied marker of oxidative tissue damage during exercise. MDA levels have been found to increase both in different tissues and plasma during exercise.^{6,8,9} It has been reported that strenuous physical exercise produces a decrease in antioxidants levels and an increase in the markers of lipid peroxidation in target tissues and blood.¹⁰ Data on changes in total antioxidant capacity (TAC) in humans are conflicting,¹¹ as is the information on the effect of exercise on lipid peroxidation.^{8,12} Evaluation of TAC is one of the most common procedures employed to evaluate the hydrosoluble antioxidant status of biological

fluids.¹³ After strenuous physical exercise, a significant increase in TAC may occur, although the opposite effect might be anticipated.¹⁴ There is ample evidence of the antioxidant activity of flavonoids both in vivo and in vitro. The results described by Ghiselli et al.¹⁴ indicate that polyphenol-rich beverages are able to transfer their antioxidant capacity to body fluids. The aim of the present study is to follow through the effects of the ingestion of a drink of polyphenolic antioxidants (flavonoids) on lipid and protein oxidation and the TAC of plasma in cyclists under oxidative stress.

Materials and methods

Subjects

A group of 31 sportsmen aged (23.6 ± 0.9 years) was selected. The subjects were cyclists, engaged in a controlled physical training program consisting of 3 h of training per day, the last week before the tests (20 h of training per week for at least 1 year). Before entry into the study, the subjects' health status was assessed from their medical history, hematological evaluation and exercise electrocardiogram. The individuals were selected consecutively among those fulfilling the inclusion criteria. The inclusion criteria involved an absence of hypertension, diabetes, cardiovascular disease, organic brain disease, alcohol or drug dependence, and any other deviation from normal food habits. The general characteristics of the participants are shown in Table 1.

All details of the study were explained to the participants, who gave their written consent. The protocol was approved by the Ethical Committee of the San Antonio Catholic University of Murcia Spain.

Dietary assessment

All volunteers were explicitly asked to follow their normal lifestyle, particularly with regard to diet and level of physical activity during the last year. A

Table 1 General characteristics of participants.

General characteristics of participants	Values
Age (years)	23.7 \pm 0.9
Weight (kg)	69.7 \pm 1.3
Height (cm)	175.7 \pm 2.4
BMI (kg/m ²)	22.9 \pm 0.8
VO _{2max} (l/min)	4.2 \pm 0.1

BMI, Body mass index; VO_{2max}, Maximal oxygen uptake. Values expressed as mean \pm SEM.

Table 2 Daily nutritional intakes of the 31 sportsmen.

	Mean	SEM
Energy intake (cal)	3188	152
Protein (g)	125	8
Carbohydrate (g)	408	21
Fat (g)	127	9
Fiber (g)	23	2
Cholesterol (mg)	255	19
Vitamin A (μ g)	2539	287
Vitamin C (mg)	133	11
Vitamin E (mg)	10	1
Zn (mg)	19	2
Mn (g)	3	0.3
Mg (mg)	390	24

Values expressed as mean \pm SEM of two 24 h surveys of each subject.

face-to-face interview was conducted with every participant and a detailed food frequency questionnaire was completed in order to obtain information about their dietary habits. The quality, quantity and frequency of consumption of red meat, chicken, fish, eggs, vegetables, fruits, dairy products and soft drinks were similar in all of the subjects. In a pre-test conducted under the direction of a qualified dietician, the subjects were instructed on how to record food intakes using standard household measures. After the food diary was completed, the dietician examined it to clarify the records and eliminate inconsistencies with the help of the subjects. Intakes were recorded twice, before each exercise-test. No statistical difference was found between the 2 day for any of the variables studied $P < 0.05$, for 7 variables: energy, protein, fat, carbohydrates, cholesterol, fiber and antioxidant vitamins (Table 2). The energy intake and dietary composition were subsequently analyzed using the AyD dietary analysis computer program.¹⁵

Analysis of antioxidant capacity of drink with polyphenols

The beverage supplied by Hero-Spain had an energy value of 42 Kcal/100 ml, and contained mainly carbohydrates from fruits concentrates 21% (black grape, raspberry and red currant), known to contain high levels of phenolic compounds;^{16–18} the beverage also contained maltodextrin, pectin, sodium citrate and vitamin C (20 mg/l), whey protein hydrolyze, sodium chloride, aroma, sweeteners

sodium cyclamate and saccharine and vitamin B1 (15% Recommended Dairy Intake—RDI).

Modal to the placebo, there was prepared by the same ingredients that the drink of study but without contributing any source of carbohydrates. His composition is basically water with sweeteners, fragrant, natural colorant, vitamin C and pectin.

The antioxidant capacity of the beverage (of the same lot of manufacture) was analyzed in three data groups, each of which was kept in darkness at different temperatures: 23, 30 and 37 °C. The measurements of the antioxidant capacity of the drink (Fig. 1) were realized for 1 year of storage, in triplicate, in a spectrophotometer Varian (mod. Cary Bio-50 UV-Vis). The capacity scavenger of radical free of the drink was evaluated by making the sample with polyphenolic antioxidants react with the radical ABTS*.¹⁹ The Trolox Equivalent Antioxidant Capacity (TEAC) assay is based on the scavenging of the 2,2'-azinobis-3-ethylbenzothiazoline-6-sulphonic acid (ABTS) radical ABTS* converting it into a colorless product. The degree of discoloration induced by a compound is related to that induced by Trolox, giving the TEAC value. Results were expressed as TEAC, defined as the millimolar mM concentration of a Trolox solution with antioxidant capacity equivalent to 1.0 mM solution of the substance under investigation.

Quantitative analysis was also carried out the concentration of total phenols of the drink was determined spectrophotometrically with the reagent Folin-Ciocalteu, using gallic acid as standard, as was qualitative analysis of the polyphenolic compounds of the drink (Table 3), by liquid chromatography in an HPLC Merck-Hitachi with L-7100 pump and using a Lichrochart 100 RP-18 reverse phase column Merck, Darmstadt, Germany (25 \times 0.4 cm, with 5 μ m particle size). For the mobile phase HPLC water with 5% of formic acid




Figure 1 Changes in antioxidant capacity (TEAC) of the replacement drink with polyphenols for a year of storage in darkness at different temperatures. Values expressed as Mean \pm SEM.

Table 3 Polyphenolic antioxidants contained in the replacement drink.

Polyphenolic antioxidants	mg/l
Anthocyanins	758.6
Flavonols	80.5
Derivates hydroxycinnamic	245.7
Stilbenes	1.0
Ellagic acid	167.7
Total	1253.5

(solvent (A) and methanol (solvent B) was used. The elution was carried out at flow speed of 1 ml/min, with increases at 30, 40 and 50 min, returning to initial conditions at 55 and allowing a settling time in the column. All the analyses were carried out in triplicate and the results are expressed as mean values \pm SEM. The different compounds were characterized by comparison chromatography following commercial models and by the visible spectrum registered with the diode net detector. The anthocyanins were quantified as cyanidin-3-rutinoside at 510 nm, the hydroxycinnamic derivatives at 320 nm as chlorogenic acid and the flavonols at 360 nm, as quercetin-3-glucoside and stilbenes as resveratrol at 320 nm (standards used).

Determination of maximal aerobic capacity

Maximal oxygen uptake was estimated by means of a continuous incremental exercise test on a electromagnetically braked cycloergometer Technogym Spin Trainer (Gambettoia, Italy), with a start load simulating a speed of 12 Km/h, increasing by 2 Km/h every minute, at a constant slope of 2%. Heart rate was continuously recorded on an electrocardiogram (ECG) G.E. Medical Systems Company; (Buc, France). Gas exchange data were obtained using an automated breath-by-breath system Sormedics Vmax Series 29-C; (Yorba Linda, CA). Relationship from the $VO_{2\max}$ —work rate and the work rate equivalent to 70% $VO_{2\max}$ was interpolated.

Exercise tests

Each subject performed two bouts of exercise on different occasions separated by one week: 1 of them testing the study drink (A), and another one with the placebo (P). The order of the trials was randomly assigned for each cyclist, but only once, before the first occasion (in this way, each cyclist play as his own control). Both beverages were

identical in appearance and taste. The flavor of the two was indistinguishable. During the exercise no other beverages or food were ingested. The trials were double-blinded (to the subjects and to the researcher).

Two 90-min rectangular tests were performed on a bicycle ergometer at 70% of the $VO_{2\max}$. Laboratory conditions were $25 \pm 0.2^\circ\text{C}$ and 70% relative humidity. The subjects drank 200 ml of beverage (A or P) every 15 min and initiated ingestion 15 minutes before the exercise. The sportsman drink 1600 ml of beverage for each test, this volume is considered usual in the habitual ingestion of liquid that the cyclists consume during the development of a sports test. Each cyclist was weighed immediately before and after exercise.

Following arrival at the laboratory at 9.00 a.m. each cyclist consumed a standard breakfast [500 ml of entire milk (Hero[®]), 200 ml of orange juice (Hero[®]), 25 g of jam of peach (Hero[®]), 2 toasts (Bimbo[®]), 1 snack of cereals (Hero Muesly[®]), sugar: 1 teaspoonful is equivalent to 3g, soluble coffee (Nescafé[®]): 1 teaspoonful is equivalent to 1g, soluble cocoa (Nesquik[®]): 1 teaspoonful is equivalent to 2g] 45 min before the beginning of the experiment. Before initiating the test each survey subject was monitored for 24 h.

Biochemical blood analyses

For each volunteer, blood samples were collected immediately before exercise, immediately after exercise and 45 min post-exercise by venous puncture from the antecubital vein. Immediately an aliquot was transferred to chilled heparinized glass tubes. Samples were centrifuged (6000 rpm/10 min), and plasma was divided into aliquots and stored at -80°C .

Measurements of hematocrit and hemoglobin

Hematocrit ratio (Htc) was measured in venous blood samples for triplicate with a microhematocrit centrifuge and corrected (0.96) for plasma trapped with the packed red cells. An additional correction (0.93) was made for the venous -to- total body Htc ratio. Hemoglobin was measured by the cyanmethemoglobin method. The methodological error for hematocrit and hemoglobin were $\pm 0.3\%$ and $\pm 0.5\%$, respectively; and percentage changes in plasma volume were estimated using the method of Dill and Costill.²⁰

Creatine kinase (CK) and lactate dehydrogenase (LDH) activity

Serum CK and LDH were used as an indirect index of exercise-induced muscle damage^{21–23} and were measured using two diagnostic kits Sigma Chemical Co. (St. Louis, MO). We have measured serum CK activity, which is used as a marker of the muscular damage, and it is associated indirectly with the increment of the permeability in the muscle cell membrane^{49,50} which it is produced in parallel way that the detection of oxidative stress induced by physical exercise. As a minimum triplicate analyses were made of each sample, and CK and LDH activities were calculated as the mean of three values that differed by no more than 10% of the lower value. The inter-assay variation coefficients (cv) were 3.5% for CK and 3.8% for LDH.

Lipid peroxidation measurement

Plasmatic levels of lipid peroxidation were measured as formation of a thiobarbituric acid (TBA) adduct of MDA, separated by HPLC.⁵¹ The sample is added to lysis buffer (0.2% Triton X-100 in $\text{KH}_2\text{PO}_4/\text{KOH}$, pH 7.4) plus butylated hydroxytoluene (100 ml of a solution of 2 g/l butylated hydroxytoluene in ethanol) (Sigma). Two 250 ml samples were incubated at room temperature for 10 min and then 500 ml 1% (w/v) TBA was added. Reaction mixtures were incubated at 90 °C for 90 min, allowed to cool and centrifuged (3000g, 10 min). Twenty microliters of the supernatant were injected into a Spherisorb ODS₂ (C18) column (Waters, Milford, MA, USA) with a guard column (Hiber C18) (Waters). Elution was performed with 65% (v/v) 50 mm $\text{KH}_2\text{PO}_4/\text{KOH}$ buffer (pH 7.0) and 35% (v/v) methanol at a flow rate of 1 ml/min. The absorbance of the sharp peak at retention time around 6 min was read at 532 nm as the level of the (TBA)₂-MDA adduct. Assays were performed in triplicate and results were expressed as $\mu\text{mol TBARS/ml}$ plasma. This method is based in the addition of some reactivities which produce cell lysis, and the products of the lipid peroxidation are poured off to the plasma and, therefore, the results are expressed as plasmatic measurements.

Plasmatic antioxidant capacity

The TAC of plasma is a quantitative indication of the state of balance of these various components in the course of time. Total antioxidant status (TAS) in plasma was measured by a chromogenic method Randox Laboratories Crumham's, North Ireland). In

this assay metmyoglobin reacts with hydrogen peroxide to form ferrylmyoglobin free radical species. Ferrylmyoglobin was incubated with the substrate 2,2'-amino-di(3-ethylbenzthiazole sulfonate) and measured at 600 nm.²⁴

Protein carbonyl measurement

Protein carbonyl content was measured by forming labelled protein hydrazone derivatives, using 2,4-dinitrophenylhydrazide (DNPH), which were then quantified spectrophotometrically.^{25–27} Briefly, after precipitation of protein with an equal volume of 1% trichloroacetic acid (TCA), the pellet was resuspended in 1 ml of DNPH 10 mM in 2 N HCl. Separate blanks were prepared by adding 1 ml of 2 N HCl without DNPH. Samples were left at room temperature for 1 h in the dark and vortexed every 15 min. An equal volume of 20% TCA was added and after centrifugation at 12,000g for 15 min at 4 °C, pellets were washed three times with 1 ml of ethanol–ethylacetate mixture (1:1) to remove the free DNPH and lipids contaminants. The final pellet was dissolved in 1 ml of 6 M urea and kept at 37 °C for 1 h in a water bath with mixer. The solution was centrifuged for 15 min at 12,000g. The carbonyl content was determined from the absorbance at 370 nm with the use of a molar absorption coefficient of 22,000 mol/l cm.

Statistical analyses

Statistical analysis was performed with the SPSS 12.0 program (Statistical Package for Social Science) for Windows. All data are presented as mean \pm SEM. Comparisons were made by Student's *t*-test for normally distributed variables, and by Wilcoxon's test for those variables that did not follow a normal distribution. The normalization adjustment for each every variable was made by means of Kolmogorov–Smirnov's test. The study of the differences between the averages at different moments was made by means of a one-way ANOVA with linear trend test. All contrasts were made considering $\alpha < 5\%$ to be an error. Correlation was evaluated using linear regression.

Results

The beverage studied showed an antioxidant capacity expressed as TEAC capacity antioxidant equivalents Trolox of 0.41 mM Trolox/ml, which does not change significantly (Fig. 1) for 6 months of storage in darkness at three different tempera-

tures (23, 30 and 37 °C) easily attainable during the transport and storage of the drink for its sale. There was a significant reduction in oxidative capacity after a year of storage at the three temperatures studied. The drink has a phenolic compound concentration of 1.43 ± 0.1 g/l, so in a test where the cyclist consumed 1600 ml, he ingested 2.3 g of polyphenolic antioxidants, mostly anthocyanins (Table 3), which have an anti-oxidative capacity equivalent to a solution of 656 mM Trolox. This antioxidant capacity is due to the polyphenols (and not to the vitamin C, antioxidant other one that forms a part of the composition of the drink), because the quantity of vitamin C contained in the drink is identical to the one that exists in the placebo. The main characteristics of the subjects are given in Table 1. The cyclists were young and some of their BMI values were within normal range. The maximum oxygen uptake, measured during cycloergometer, showed also that the subjects' VO_2 values were well adapted to the physical activity that the cyclists in the study reported. "Regional and national level" is referred to the category, that is where the competition take place, not in relation with the level of training. The maximum oxygen uptake, measured during cycloergometer, showed also that the subjects' $VO_{2\max}$ values were well adapted to the level of training that the cyclists in the study reported (20 h/week). The data of $VO_{2\max}$ in relation to body weight is $VO_{2\max} = 58.9 \pm 4.2$ (ml min kg)⁻¹.

The analysis of nutritional status (Table 2) showed that energy intake was similar in the two surveys made of each cyclist (3188 cal/day). Intakes were recorded twice, before each exercise-test:

Firstly, 7 days before (without any dietary orders).

Secondly, the day before the exercise test (with the same dietary orders, in reference to avoid the foods rich in polyphenols: tea, coffee, red grapes, wine, red fruits, etc). The same dietary orders were given to the volunteers the last 3 days before the tests.

The data shown in Table 2 make it clear that the subjects consume balanced diets in terms of macronutrient composition. The means \pm SEM percentages of total dietary energy intake from carbohydrates, protein and fat was $50 \pm 5\%$, $14 \pm 1\%$ and $36 \pm 4\%$, respectively. Observed differences of nutrient intake were not statistically significant.

To correct for changes in plasma volume that might occur over the course of the exercise-tests, hemoglobin levels and hematocrit were measured both before and after each trial. Mean levels of plasma hemoglobin and hematocrit remained unchanged, and the mean percentage change in plasma volume was small (Table 4), but individual percentage changes ranged from -5.1% to $+6.1\%$, making individual correction of plasma values important.

The CK values (Table 5) coincides with the results obtained by Hartman et al.,²⁸ who observed that sportsmen with chronically high base CK levels show great variability. CK values increased significantly after exercise, both in the placebo test ($p < 0.05$) and when the cyclists consumed the drink with polyphenols ($p < 0.05$), which indicates that the physical activity of the test caused muscular damage to the cyclists. The theory that muscular damage is responsible for the post-exercise inflammatory stress reaction is based on the following findings: immediately after exercise-induced muscle damage, neutrophils infiltrate the muscle, and begin the removal of degraded proteins and cell debris, followed by monocytes and macrophages. These cells are capable of producing ROS and secreting pro-inflammatory cytokines such as IL-1, TNF- α or IL-8, leading to oxidative stress and eventually overshooting inflammation.²⁹

LDH values observed before (I) and after the exercise (F) (Table 5) were higher than the normal values of the enzyme in adults 140–280 U/l. Nevertheless significant differences were not found though there was a trend to increase between LDH values after the test (F) and before exercise (I). These results coincide with those found by

Table 4 Values of hematocrit (Hc), hemoglobin (Hb) and changes in plasmatic volume (% Δ PV) pre- and post-exercise ($n = 31$).

Variable	AI	AF	PI	PF
Hb (g/100 ml)	15.3 ± 0.2	15.3 ± 0.2	15.2 ± 0.2	15.4 ± 0.2
Htc \blacklozenge (%)	44 ± 0.4	44 ± 0.4	44 ± 0.5	45 ± 0.5
% Δ PV		-0.1 ± 0.9		-2.1 ± 0.7

Values expressed as mean \pm SEM; Htc \blacklozenge = Htc \times 0.96; A: Supplemented test with polyphenolic antioxidants), P: Placebo test; I: Pre-exercise, F: Post-exercise (45 min post-exercise).

Anthony et al.³⁰ The fact that significant differences were not observed before and after the exercise may be due to the differences found among individuals, which justifies the inclusion in this type of studies of both tests placebo and with the drink under study on the same individual. At rest, the TBARS values (I) in two test (A) and (P) did not show significant differences (Fig. 2). This indicates the homogeneity of the population before the tests. In the placebo test the TBARS value immediately after exercise (PFo) is significantly ($P < 0.05$) above the value before exercise (PI), which allows us to affirm that the physical activity of the cyclists has induced oxidative stress. In the test in which the cyclist consumes the drink with polyphenols the value of TBARS immediately after exercise (AFo) does not increase significantly. TBARS concentration in both tests showed a significant increase immediately after strenuous exercise (Fo), whereas 45 min after the tests (F) TBARS concentration diminished and the pre-exercise values (I) were recovered (Fig. 2). These results coincide with those of other authors.^{31,32}

TAC is considered a valuable measurement of the antioxidant status of biological fluids and tissues.²⁴ The TAS values obtained in cyclists pre- and post-exercise were lower than the adult (reference values 1.30–1.77 nmol/l plasma). As can be seen in Fig. 3, no significant differences were found in TAS plasmatic pre- and post-exercise in the two tests (F vs. I), even when the placebo and the supplemented drink were compared. Polyphenolic antioxidants supplementation had no effect on plasmatic antioxidant capacity levels in the cyclists of this study.




Figure 2 Thiobarbituric acid-reactive substances (TBARS; by-products of lipid peroxidation) in blood plasma before (I) (black bars), immediately after (fo), (ruled bars), and 45 min after (F), (white bars) exercise (polyphenol-supplemented (A) and placebo (P)). Values expressed as mean \pm SEM. *Significant difference ($p < 0.05$) fo vs. I.

Protein oxidized, expressed as nmol of carbonyl groups/mg of protein, did not significantly differ at pre-exercise among the two tests. The mean values were: for test A, 1.02 ± 0.12 in pre-exercise, 0.78 ± 0.09 in post-exercise; and for test P, 0.90 ± 0.10 in pre-exercise, 1.02 ± 0.13 in post-exercise (Fig. 4). Polyphenolic antioxidant supplementation led to a




Figure 3 Antioxidant capacity in blood plasma before (I), (solid bars), and 45 min after (F), (crosshatched bars) exercise (polyphenol-supplemented (A) and placebo (P)). Values are means \pm SEM of total antioxidant status (TAS).




Figure 4 Protein oxidation in blood plasma before (I), (solid bars), and 45 min after (F), (crosshatched bars) exercise (polyphenol-supplemented (A) and placebo (P)). Values are means \pm SEM of carbonyl groups/mg of plasmatic protein. *Significant difference ($p < 0.05$) F vs. I. **Significant difference ($p < 0.05$) A vs. P.

Table 5 Descriptive CK and LDH (U/l) statistics pre- and post-exercise for both tests (A and P).

Description (n = 31)	Mean	SEM
CK AI (U/l)	203.0	7.4
CK AF (U/l)	233.1*	8.4
CK PI (U/l)	128.4	2.9
CK PF (U/l)	168.7*	3.8
LDH AI (U/l)	329.2	3.8
LDH AF (U/l)	337.4	6.8
LDH PI (U/l)	331.5	3.9
LDH PF (U/l)	349.1	3.2

Values corrected for plasma volume. A: Antioxidant supplement test, P: Placebo Test, I: Pre-exercise, F: Post-exercise. *Significant increase ($p < 0.05$) F vs. I.

23% decrease in mean protein carbonyl levels ($p < 0.05$), but the placebo test showed an increase of 12% in carbonyl proteins. These results coincide with those of O'Byrne et al.³³ who found a decrease of 20% in carbonyl groups levels after supplementation of subjects with grape juice as a source of flavonoids. Alessio et al.³⁴ compare the "pre-post" increases in the protein oxidation carbonyl groups between stressful aerobic exercise and other isometric anaerobic exercise, finding increases of 67% and 12%, respectively.

Discussion

This work presents the results from a controlled experiment where the effect of polyphenolic antioxidants on changes in plasma lipid peroxidation, capacity antioxidant and protein oxidation, were investigated in trained cyclists. The antioxidant effect of the beverage could be due to polyphenolic content and not to the vitamin C (another antioxidant component of the drink) due to the fact that the placebo used in the study contains the same quantity of vitamin C that the designed drink, and the statistical analysis is realized on the basis of the opposing differences drink vs. placebo, in the different biomarkers of oxidative stress analyzed. A group of young cyclists engaged in a controlled physical training programme showed an improvement in the plasmatic oxidative stress biomarkers after ingestion of a beverage with polyphenolic antioxidants. In this group of trained subjects, a significant increase in CK and TBARS levels was found in relation to exercise, which suggests that sub-maximal physical activity 90 min at 70% of VO_2 on the cycloergometer produces muscular damage and oxidative stress.³⁷ Under these conditions in the organism the utilization of fat as energetic fuel increases, but this comes neither from the oily plasmatic acids nor from the adipose tissue reserves; rather the lipid substratum used as energetic fuel is intramuscular fat.³⁸ For this reason, in the study of trained sportsmen, there is justifiably a significant increase (5.6%) in lipid peroxidation. Nevertheless, in the polyphenolic antioxidant supplement test no significant increase in lipoperoxidation was observed. Exercise-associated increases in secondary products of lipid peroxidation in plasma have often been used as a marker of oxidative stress.³⁹⁻⁴¹ It is critically important to stabilize lipid peroxidation by products in biological samples by immediate treatment with BHT, because lipid peroxides are not only markers of oxidative stress but are also

strong pro-oxidants themselves. Lipid peroxy radicals possess sufficient energy to initiate further lipid peroxidation chain reactions.

Flavonoids and cinnamic acids the main polyphenols in the drink are known as primary antioxidants^{35,42} and act as free radical acceptors and chain breakers. The ability of flavonoids to form complexes with a cupric ion has also been demonstrated by UV spectral studies.⁴³ Such complexities may contribute to the antioxidative action of flavonoids. Bioavailability studies have also shown that the concentrations of intact flavonoids in human plasma rarely exceed $1 \mu\text{M}$ when the quantities of polyphenols ingested do not exceed those commonly ingested with our diets, whereas the cyclists in this study ingest a quantity greater than those commonly ingested in our diet. These maximum concentrations are most often reached 1-2 h after ingestion.⁴⁴ For most flavonoids absorbed in the small intestine, the plasmatic concentration rapidly decreases (half-life period of 1-2 h).⁴⁵ The limited data available indicate that concentrations are potentially high enough to produce the biological effects observed in this study.

Having carried out sub-maximal physical activity for 90 min about 14% of the energy obtained by the body comes from protein oxidation,⁴⁶ and in addition it is necessary to consider the existence of high protein turnover, due not only to the alanine cycle but also to the muscular micro-injury induced by exercise. Nevertheless the fact that the study sample consists of trained sportsmen, and that the lipid substratum is more sensitive than the protein substratum to oxidation, justifies the fact that after placebo test the results did not show a significant increase in oxidized protein. A significant 23% of decrease was observed in plasma protein carbonyl levels after the polyphenol supplement test.

In our study the results did not show significant variations in the antioxidant plasmatic capacity of the cyclists after the test with regard to the values obtained before initiating the test. Neither was significant differences found in the antioxidant plasmatic capacity of cyclists as a consequence of the ingestion of the polyphenolic antioxidant drink. During strenuous physical activity the level of uric acid increases,⁴⁷ contributing 19.3% to the total antioxidant plasmatic capacity, as opposed to the scant contribution of exogenous antioxidants such as ascorbic acid (3.1%) and α -tocopherol (1.7%).⁴⁵ All this suggests that the polyphenolic compounds responsible for the antioxidant capacity of the drink consumed by the cyclists in our study, in spite of having a greater antioxidant capacity than ascorbic acid and α -tocopherol, his scanty bioavail-

ability and therefore his scanty contribution to the antioxidant plasmatic capacity is not sufficient to modify the whole antioxidant plasmatic state whole in the studied sportsmen. Nevertheless, the results observed in the decrease of the levels of carbonyl groups (as markers of protein oxidation) in the cyclists when they consume the polyphenol drink, in contrast to the same cyclists when they consume a placebo, indicate a decrease in the oxidative damage induced by physical activity in spite of the total antioxidant state in the plasma remaining the same. Rossi et al.⁴⁸ have demonstrated an improvement in the antioxidant state and a decrease in the oxidative damage induced by exercise with the ingestion of soybean, and suggest that this improvement is attributable to the isoflavones (other flavonoids) of the soybean.

Statistical analysis of the results obtained during this study in different biomarkers of oxidative stress in plasma lipid peroxidation, protein oxidation, creatinquinase and LDH enzymes, and total antioxidant state, shows that the analysis of protein oxidation is the biomarker that in the study conditions reflects the appearance of oxidative stress as a consequence of physical activity on the cycloergometer. Similarly, the variations observed amongst individuals in all the biomarkers suggest that in this type of study it is crucial to use the same population sample as a control group.

Conclusions

In trained cyclists, aerobic physical activity in the effort tests in the laboratory for 90 min at 70% VO_2 max generated muscular damage and oxidative stress caused by exercise.

Compared to a placebo, the drink with polyphenolic antioxidants proved capable of reducing the degree of protein oxidation caused by physical activity. In this drink the antioxidant capacity is kept constant for 6 months, which allows it to be marketed as a replacement drink for sportsmen.

The high variations observed among individuals in the biomarkers analyzed serves to support the idea that in order to study the effect of antioxidant on oxidative stress it is necessary to use as a control group the same subjects that receive the antioxidant supplement.

Acknowledgments

Thanks are due to the sportsmen, trainers and managers of the cycling clubs who participated. We

are also grateful to Professors A. Luque and A.B. Martínez for their personal work, and to Gabriel, Sergio and Almudena for their technical assistance.

References

1. Bors W, Michel C, Stettmaier K. Antioxidant effects of flavonoids. *Biofactors* 1997;64:399–402.
2. Van Acker FA, Schouten O, Haenen GR, Van der Vijgh WJ, Bast A. Flavonoids can replace alpha-tocopherol as an antioxidant. *FEBS Lett* 2000;473:145–8.
3. Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr Rev* 1998;56(11):317–33.
4. Lee MJ, Wang ZY, Li H, et al. Analysis of plasma and urinary tea polyphenols in human subjects. *Cancer Epidemiol Biomarkers Prev* 1995;4(4):393–9.
5. Unno T, Kondo K, Itakura H, Takeo T. Analysis of (–)-epigallocatechin gallate in human serum obtained after ingesting green tea. *Biosci Biotechnol Biochem* 1996;60:2066–8.
6. Alessio H. Exercise-induced oxidative stress. *Med Sci Sports Exerc* 1993;23:218–24.
7. Ji LL. Exercise and oxidative stress: role of the cellular antioxidant systems. *Free Rad Biol Med* 1994;16:339–46.
8. Duthie GG, Robertson JD, Maughan RJ, Morrice PC. Blood antioxidant status and erythrocyte lipid peroxidation following distance running. *Arch Biochem Biophys* 1990;282(1):78–83.
9. Marzatico F, Pansarasa O, Bertorelli L, Somenzini L, Della Valle G. Blood free radical antioxidant enzymes and lipid peroxides following long-distance and lactacidemic performances in highly trained aerobic and sprint athletes. *J Sports Med Phys Fitness* 1997;37(4):235–9.
10. Davies KJA, Quintanilha AT, Brooks GA, Packer L. Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun* 1982;107(4):1198–205.
11. Ohno H, Sato Y, Yamashita K, Doi R, Arai K, Kondo T. The effect of brief physical exercise on free radical scavenging enzyme systems in human red blood cells. *Can J Physiol Pharmacol* 1986;64:1263–5.
12. Viguie CA, Frei B, Shigenaga MK, Ames BN, Packer L, Brooks GA. Antioxidant status and indexes of oxidative stress during consecutive days of exercise. *J Appl Physiol* 1993;75:566–72.
13. Brites FD, Evelson PA, Christiansen MG, et al. Soccer players under regular training show oxidative stress but an improved plasma antioxidant status. *Clin Sci* 1999;96:381–5.
14. Ghiselli A, Serafini M, Natella F, Scaccini C. Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Radic Biol Med* 2000;29(11):1106–14.
15. Villegas García JA. Alimentación y Salud (programa informático), Versión 4.2. 1996.
16. Alcolea JF, Cano A, Acosta M, Arnao MB. Hydrophilic and lipophilic antioxidant activities of grapes. *Nahrung* 2002;46(5):353–6.
17. Liu M, Li XQ, Weber C, Lee CY, Brown J, Liu RH. Antioxidant and antiproliferative activities of raspberries. *J Agric Food Chem* 2002;50(10):2926–30.
18. Maatta K, Kamal-Eldin A, Torronen R. Phenolic compounds in berries of black, red, green, and white currants (*Ribes* sp.). *Antioxid Redox Signal* 2001;3(6):981–93.

19. Espín JC, Wichers HJ. Study of the oxidation of resveratrol catalyzed by polyphenol oxidase. Effect of polyphenol oxidase, lacase and peroxidase on the antiradical capacity of resveratrol. *J Food Biochem* 2000;24:225–50.
20. Dill DB, Costill DL. Calculation of percentage changes in volumes of blood, plasma, and red cells in dehydration. *J Appl Physiol* 1974;37:247–8.
21. Agarwal MM, Ankra-Badu GA. Serum creatine kinase: a marker for muscle damage in sickle cell painful crisis. *Ann Saudi Med* 1999;19(3):264–6.
22. Chung MS, Park H, Chang WH. The values of serum creatine kinase and lactate dehydrogenase in Korean Olympic athletes during training season. In: Proceedings of the fifth IOC congress on sport sciences with the annual conference of science and medicine in sports, 1999.
23. Ji LL. Antioxidants and oxidative stress in exercise. *Proc Soc Exp Biol Med* 2000;222:283–92.
24. Miller NJ, Rice-Evans C, Davies ML, Gopinathan V, Milner A. A novel method for measuring antioxidant capacity and its applications to monitoring the antioxidant status in premature neonates. *Clin Sci* 1993;84:407–12.
25. Evans P, Lyras L, Halliwell B. Measurement of protein carbonyls in human brain tissue. *Methods Enzymol* 1999;300:145–56.
26. Levine RL, Williams JA, Stadman ER, Shacter E. Carbonyls assays for determination of oxidatively modified proteins. *Methods Enzymol* 1994;233:346–57.
27. Reznick AZ, Packer L. Oxidative damage to proteins: spectrophotometric method for carbonyl assay. *Methods Enzymol* 1994;233:357–63.
28. Hartman U, Mester J. Training and overtraining markers in selected sports. *J Sci Sport Exerc* 2000;32(1):209–15.
29. Ji LL. Antioxidants and Oxidative Stress in Exercise. *Proc Soc Exp Biol Med* 2000;222:283–92.
30. Anthony ME, Sparks K, Little K, Rood A, Vangelos Z. (on line) (access 11 de enero de 2003) URL disponible en: <http://www.run-insight.com/RI-SS-FinalMarathonStudy.doc>
31. Krotkiewski M, Brzezinska Z. Lipid peroxides production after strenuous exercise and in relation to muscle morphology and capillarization. *Muscle Nerve* 1996;19(12):1530–7.
32. Lovlin R, Cottle W, Pyke L, Kavanagh M, Belcastro AN. Are indices of free radical damage related to exercise intensity? *Eur J Appl Physiol* 1987;56(3):313–6.
33. O'Byrne DJ, Devaraj S, Grundy SM, Jialal I. Comparison of the antioxidant effects of Concord grape juice flavonoids and alpha-tocopherol on markers of oxidative stress in healthy adults. *Am J Clin Nutr* 2002;76(6):1367–74.
34. Alessio HM, Hagerman AE, Fulkerson BK, Ambrose J, Rice RE, Wiley RL. Generation of reactive oxygen species after exhaustive aerobic and isometric exercise. *Med Sci Sports Exerc* 2000;32(9):1576–81.
35. Rice Evans CA, Miller NJ, Paganga G. Antioxidant properties of phenolics compounds. *Trends Plant Sci* 1997;2:152–9.
37. König D, Berg A. Exercise and oxidative stress: is there a need for additional antioxidants. *Öster J Sport Med* 2002;3(1):6–15.
38. Córdova A. *Fisiología Dinámica*. Barcelona: Masson; 2003.
39. Jenkins RR, Krause K, Schofield LS. Influence of exercise on clearance of oxidant stress products and loosely bound iron. *Med Sci Sports Exerc* 1993;25:213–37.
40. Ji LL, Katz A, Fu R, Griffiths M, Spencer M. Blood glutathione status during exercise: effect of carbohydrate supplementation. *J Appl Physiol* 1993;74:788–92.
41. Kretzschmar M, Müller D, Hübscher J, Marin E, Klinger W. Influence of aging, training and acute physical exercise on plasma glutathione and lipid peroxides. *Int J Sports Med* 1991;12:218–22.
42. Pratt DE, Hudson B. In: Hudson B, editor. *Food antioxidants*. Amsterdam: Elsevier; 1990. p. 171–91.
43. Hudson B, Lewis JI. Polyhydroxy flavanoids—antioxidants for edible oils. *Food Chem* 1983;14:45–81.
44. Lee MJ, Wang ZY, Li H, et al. Analysis of plasma and urinary tea polyphenols in human subjects. *Cancer Epidemiol Biomarkers Prev* 1995;4:393–9.
45. Lapidot T, Harel S, Granit R, Kanner J. Bioavailability of red wine anthocyanins as detected in human urine. *J Agric Food Chem* 1998;46:4297–302.
46. Costill D, Sherman WM, Fink WJ. The role of dietary carbohydrates in muscle glycogen resynthesis after strenuous running. *Am J Clin Nutr* 1981;34:831–6.
47. Gómez-Cabrera MC, Gimeno A, Lloret A, Miñana JB, Marquez R, Viña J. Deporte de alta competición y daño oxidativo: papel de los nutrientes antioxidantes. (on line)(access 21 de mayo de 2001). URL: www.antioxidantes.com.ar/12/Art118.htm
48. Rossi AL, Blostein-Fujii A, DiSilvestro RA. Soy beverage consumption by young men: increased plasma total antioxidant status and decreased acute, exercise-induced muscle damage. *J Nutr Func Med Foods* 2000;3(1):279–91.
49. Van der Meulen JH, Kuipers H, Drukker J. Relationship between exercise-induced muscle damage and enzyme release in rats. *J Appl Physiol* 1991;71:999–1004.
50. Child R, Brown S, Day S, Donnelly A, Roper H, Saxton J. Changes in indices of antioxidant status, lipid peroxidation and inflammation in human skeletal muscle after eccentric muscle actions. *Clin Sci* 1999;96:105–15.
51. Chirico S. High-performance liquid chromatography-based thiobarbituric acid tests. *Methods Enzymol* 1994;223:314–8.

Further Reading

36. Song J, Kwon O, Chen S. Flavonoid inhibition of sodium-dependent vitamin C transporter 1 (SVCT1) and glucose transporter isoform 2 (GLUT2), intestinal transporters for vitamin C and glucose. *J Biol Chem* 2002;277(18):15252–60.

Available online at www.sciencedirect.com

